

## Musculoskeletal System NCLEX Review

### Musculoskeletal Disorders

#### Diagnostic Tests

**Radiography and magnetic resonance imaging (MRI)** are commonly used procedures to diagnose disorders of the musculoskeletal system.

#### Nursing Interventions:

- Handle injured areas carefully and support extremities above and below the joint.
- Remove any radiopaque and metallic objects, such as jewelry.
- Shield the client's testes, ovaries, or pregnant abdomen.
- **NO** to pregnant or expecting mothers.
- Inform that the radiation from radiography is minimal and not dangerous.
- *\*Administer analgesics as prescribed **before** the procedure.*

**Arthrocentesis** is used to diagnose joint inflammation and infection; involves aspirating synovial fluid, blood, or pus via a needle inserted into a joint cavity.

#### Nursing Interventions:

- Obtain informed consent
- Apply an elastic compression bandage post procedure
- Use ice to decrease pain and swelling.
- Administer analgesics as prescribed.
- Pain can continue for up to 2 days after administration of corticosteroids into a joint.
- Instruct the client to rest the joint for 8 to 24 hours post procedure.

**Arthroscopy** provides an endoscopic examination of various joints; articular cartilage abnormalities can be assessed, loose bodies removed, and the cartilage trimmed.

#### Nursing Interventions:

- Fasting: 8 to 12 hours before the procedure.
- Obtain informed consent.
- Assess the neurovascular status of the affected extremity.
- An elastic compression bandage should be worn post procedure for 2 to 4 days.
- Instruct the client that walking with weight bearing usually is permitted after sensation returns but to limit activity for 1 to 4 days.
- Instruct the client to elevate the extremity as often as possible for 24 hours post-op.
- Place ice on the site.

**Bone scan** is used to identify, evaluate, and stage bone cancer before and after treatment; it is also used to detect fractures.

## Nursing Interventions:

- **NPO** prior to the procedure.
- Obtain informed consent.
- Remove all jewelry and metal objects.
- *Increase OFI* after the procedure to eliminate excess isotopes
- 1 to 3 hours after the injection, have the client void before the scanning procedure is completed.

**Bone or muscle biopsy** may be done during surgery or through aspiration or punch or needle biopsy.

## Nursing Interventions:

- Obtain informed consent
- Monitor for bleeding, swelling, hematoma, or severe pain.
- Elevate the site for 24 hours following the procedure.
- Apply ice packs.
- Monitor for signs of infection.

**Electromyography (EMG)** measures electrical potential associated with skeletal muscle contractions.

## Nursing Interventions:

- Obtain informed consent.
- Instruct the client that the needle insertion is uncomfortable.
- Do not take any stimulants or sedatives for 24 hours before the procedure.
- Inform the client that slight bruising may occur at the needle insertion sites.

## Injuries

**Strains** are an excessive stretching of a muscle or tendon.

## Nursing Interventions:

- Apply warm and cold compress
- Limit activity
- Medicate for pain and muscle relaxants

**Sprains** are characterized by pain and swelling.

## Nursing Interventions:

### " R-I-C-E "

- **R**est the affected site
- Apply **I**ce compress
- Apply **C**ompression bandage

- Elevate legs
- \*Casting may require

**Rotator cuff injuries** is characterized by shoulder pain and the inability to maintain abduction of the arm at the shoulder (drop arm test).

### **Nursing Interventions:**

- Apply **ice to heat compress**.
- Apply arm sling.
- Administer NSAIDs as prescribed.

**Fractures** a break in the continuity of the bone caused by trauma, twisting because of muscle spasm or indirect loss of leverage, or bone decalcification and disease.

### **Signs and Symptoms:**

#### **" BROKEN "**

- **Bruising** with pain and swelling
- **Reduced movement**
- **Odd appearance**
- **Kracking sounds**
- **Edema and erythema on sites**
- **Neurovascular impairment** (decrease in sensation, temperature changes, loss of function etc.)

### **Types of Fractures:**

**Closed or Simple:** Skin over the fractured area remains intact.

**Comminuted:** The bone is splintered or crushed, creating numerous fragments.

**Complete:** The bone is separated completely by a break into 2 parts.

**Compression:** A fractured bone is compressed by other bone.

**Depressed:** Bone fragments are driven inward.

**Greenstick:** One side of the bone is broken and the other is bent; these fractures occur most **commonly in children**.

**Impacted:** A part of the fractured bone is driven into another bone.

**Incomplete:** Fracture line does not extend through the full transverse width of the bone.

**Oblique:** The fracture line runs at an angle across the axis of the bone.

**Open or Compound:** The bone is exposed to air through a break in the skin, and soft tissue injury

and infection are common.

**Pathological:** The fracture results from weakening of the bone structure by pathological processes such as neoplasia; also called spontaneous fracture.

**Spiral:** The break partially encircles bone. **Transverse:** The bone is fractured straight across.

### Nursing Interventions:

- *Immobilize* the affected extremity with a cast or splint.
- *Assess the neurovascular status* of the extremity.
- Interventions for a fracture: Reduction, fixation, traction, cast

**Traction** is the exertion of a pulling force applied in 2 directions to reduce and immobilize a fracture; provides proper bone alignment and reduces muscle spasms.

### Nursing Interventions:

#### " TRACTION "

- Temperature (infection, extremity)
- Ropes hang freely
- Alignments
- Circulation check
- Type and location of fracture
- Increase OFI
- Overhead trapeze
- No weight on bed or floor

### Types of tractions:


**Skeletal traction** is applied mechanically to the bone with pins, wires, or tongs; typical weight for skeletal traction is 25 to 40 lb (11 to 18 kg).

### Nursing Interventions:


- Monitor color, motion, and sensation of the affected extremity.
- Monitor the insertion sites for redness, swelling, drainage, or increased pain.
- Provide insertion site care as prescribed.

**Skin traction** is applied by using elastic bandages or adhesive, foam boot, or sling.

## Skin Traction




**Cervical Skin Traction** relieves muscle spasms and compression in the upper extremities and neck.



## Nursing Interventions:

- Uses ahead halter and chin pad to attach the traction.
- Use **powder** to protect the ears from friction rub.
- Elevate the head of the bed **30 to 40 degrees**.
- Attach the weights to a pulley system over the head of the bed.

**Buck's(extension) skin traction** is used to alleviate muscle spasms and immobilize a lower limb by maintaining a straight pull on the limb with the use of weights.



## Nursing Interventions:


- Apply boot appliance to the attach to the traction.
- Not more than 8 to 10 lb (3.5 to 4.5 kg) of weight should be applied as prescribed.
- Elevate the foot of the bed to provide the traction.

**Pelvic Skin Traction** is used to relieve low back, hip, or leg pain or to reduce muscle spasm.

### Nursing Interventions:

- Apply the traction belt snugly over the pelvis and iliac crest and attach to the weights.
- Use measures as prescribed to prevent the client from slipping down in bed.

**Balanced Suspension Traction** Used to approximate fractures of the femur, tibia, or fibula; produced by a counterforce other than the client.



### Nursing Interventions:

- Position the client in a **low Fowler's position** on either the side or the back.
- Maintain a **20-degree angle** from the thigh to the bed.
- Protect the skin from breakdown.
- Provide pin care (if pins are used with the skeletal traction)
- Clean the pin sites with sterile normal saline and hydrogen peroxide or povidone-iodine.

**Casts** are used to immobilize bones and joints into correct alignment after a fracture or injury using plaster, fiberglass, or air casts.

### Nursing Interventions:

- Keep the cast and extremity elevated.
- Allow a wet plaster cast **24 to 72 hours** to dry (synthetic casts dry in 20 minutes).
- Handle a wet plaster cast with the palms of the hands (not fingertips) until dry.
- Turn the extremity every **1 to 2 hours**, unless contraindicated.
- A hair dryer can be used on a cool setting to dry
- Monitor closely for circulatory impairment (compartment syndrome)
- Maintain smooth edges around the cast to prevent crumbling of the cast material.
- **Monitor for signs of infection** such as increased temperature, hot spots on the cast, foul odor, or changes in pain.

- Instruct the client not to stick objects inside the cast.
- Teach the client to keep the cast clean and dry.
- Instruct the client in isometric exercises to prevent muscle atrophy.

**Caution!** Monitor a casted extremity for circulatory impairment such as pain, swelling, discoloration, tingling, numbness, coolness, or diminished pulse. Notify the HCP immediately if circulatory compromise occurs.

## Complications of Fractures:

### Fat Embolism

- Notify the health care provider (HCP).
- Administer oxygen.
- 3. Administer intravenous (IV) fluids
- Monitor vital signs and respiratory status.
- Prepare for intubation and mechanical ventilation if necessary
- Follow up on results of diagnostic tests such as chest x-ray or CT scan
- Documentation

**Pulmonary Embolism** is caused by the movement of foreign particles (blood clot, fat, or air) into the pulmonary circulation.

### Signs and Symptoms:

- Restlessness and apprehension
- Sudden onset of dyspnea and chest pain
- Cough, hemoptysis, hypoxemia, or crackles

### Nursing Interventions:

- Notify the HCP immediately
- Administer oxygen, intravenous (IV), anticoagulant therapy.

**Compartment Syndrome** occurs when pressure increases within 1 or more compartments, leading to decreased blood flow, tissue ischemia, and neurovascular impairment.

**Note:** \* *Neurovascular damage may be irreversible if not treated within 4 to 6 hours.*

### Signs and Symptoms:

#### " 5Ps "

- **P**ain (increasing) in the limb.
- **P**ale, dusky, or edematous distal tissues.
- **P**ain with passive movement
- **P**aresthesia
- **P**ulselessness (*late sign*)

## Nursing Interventions:

- Notify the HCP immediately
- Elevate the affected extremity.
- Fasciotomy to relieve pressure and restore tissue perfusion.
- Loosen tight dressings or bivalve restrictive cast as prescribed.

**Osteomyelitis** (inflammatory response in bone tissue) can be caused by the introduction of organisms into bones leading to localized bone infection.

## Signs and Symptoms:

- Tachycardia and fever
- Erythema and pain in the area
- Leukocytosis and elevated ESR level

## Nursing Interventions:

- Notify the HCP
- Prepare to initiate aggressive, long-term IV antibiotic therapy.

**Avascular necrosis** occurs when a fracture interrupts the blood supply to a section of bone, leading to bone death.

## Signs and Symptoms:

- Pain
- Decreased sensation

## Nursing Interventions:

- Notify the HCP if pain or numbness occurs.
- Prepare the client for removal of necrotic tissue.

## Crutch Walking

*\*This video includes all the things you need to learn regarding the use of crutches.*

*Please click the " Link " below to watch this video.*

[Link](#)

## Nursing Interventions:

- The distance between the axillae and the arm: **2 to 3 finger widths** in the axilla space.
- The elbows should be slightly flexed, **20 to 30 degrees**, when the client is walking.
- When ambulating with the client, **stand on the affected side**.
- Look up and outward when ambulating and to place the crutches **6 to 10 inches (25.5 cm)** diagonally in front of the foot.

- Instruct the client to stop ambulation if numbness or tingling in the hands or arms occurs.

**Hemicanes or quadripod canes** are used for clients who have the use of only 1 upper extremity.



### **Nursing Interventions:**

- Position the cane at the client's unaffected side, with the straight, nonangled side adjacent to the body.
- Position the cane **6 inches (15 cm)** from the unaffected client's side.

### **Walker**

### **Nursing Interventions:**

- Stand adjacent to the client on the affected side.
- Instruct the client to put all 4 points of the walker flat on the floor before putting weight on the hand pieces.
- Instruct the client to move the walker forward, followed by the affected or weaker foot and then the unaffected foot.

### **Fractured Hip**

### **Types:**

1. **Intracapsular** (femoral head is broken within the joint capsule); Femoral head and neck receive decreased blood supply and heal slowly.

### Nursing Interventions:

- Skin traction is applied preoperatively to reduce the fracture and decrease muscle spasms
- *Treatment:* Total hip replacement or open reduction internal fixation (ORIF) with femoral head replacement.
- To prevent hip displacement postoperatively; avoid extreme hip flexion.

1. **Extracapsular** (fracture is outside the joint capsule) fracture can occur at the greater trochanter or can be an intertrochanteric fracture.

### Nursing Interventions:

- *Preoperative treatment:* Balanced suspension or skin traction
- *Surgical treatment:* ORIF with nail plate, screws, pins, or wires.
- *Postoperative treatment:* Monitor for signs of delirium and institute safety measures. Prevent internal or external rotation; avoid extreme hip flexion.
- Elevate the head of the bed 30 to 45 degrees for **meals only**.
- Avoid weight-bearing on the affected leg.
- Keep the operative leg extended, supported, and elevated (preventing hip flexion)
- Monitor for wound infection or hemorrhage.
- Use antiembolism stockings or sequential compression stockings.
- Avoid crossing the legs and activities that require bending over.

**Total knee replacement** is the implantation of a device to substitute for the femoral condyles and tibial joint surfaces.

### Post-Op Care:

- Monitor surgical incision for drainage and infection.
- Prepare the client for out-of-bed activities as prescribed; **avoid leg dangling**.
- Weight-bearing with an assistive device is prescribed as tolerated.
- Administer antibiotics if prescribed.

**Joint Dislocation** injury of the ligaments surrounding a joint, which leads to displacement or separating of the articular surfaces of the joint.

**Subluxation** incomplete displacement of joint surfaces when forces disrupt the soft tissue that surrounds the joints.

### Signs and Symptoms:

- Asymmetry of the contour of affected body parts.
- Pain, tenderness, dysfunction, and swelling
- X-rays are taken to determine joint shifting.

### Nursing Interventions:

- **Focus of treatment:** pain relief, joint support, and joint protection.
- Open or closed reduction is done with a postprocedural joint immobilization.
- Initial activity restriction is followed by gentle range-of-motion activities and a gradual return of activities to normal levels.

**Cervical Disk Herniation** occurs at the **C5 to C6** and **C6 to C7** interspaces; causes pain radiation to shoulders, arms, hands, scapulae, and pectoral muscles.

### Signs and Symptoms:

- Paresthesia
- Numbness
- Weakness of the upper extremities

### Nursing Interventions:

- Bed rest
- Immobilize the cervical area with a cervical collar or brace.
- Apply **heat** to reduce muscle spasms and apply **ice** to reduce inflammation and swelling.
- Maintain head and spine alignment.
- Administer analgesics, corticosteroids, sedatives, and anti-inflammatory medications as prescribed.
- Avoid flexing, extending, and rotating the neck.
- Avoid the prone position; maintain in neutral position

**Lumbar Disk Herniation** most often occurs at the L4 to L5 or L5 to S1 interspace. Pain is relieved by bed rest and aggravated by movement, lifting, straining, and coughing.

### Signs and Symptoms:

- Lower back pain radiating to lower limbs down.
- Muscle spasm to lower extremities.

### Nursing Interventions:

- Apply **heat** to decrease muscle spasms and apply **ice** to decrease inflammation and swelling.
- Instruct the client to sleep on the side, with the knees and hips flexed, and place a pillow between the legs.
- Apply pelvic traction as prescribed to relieve muscle spasms and decrease pain.
- Begin progressive ambulation as inflammation, edema, and pain subside.
- Instruct the client about application techniques for corsets or braces to maintain immobilization and proper spine alignment.
- Instruct about proper body mechanics.

**Disk Surgery** is used when spinal cord compression is suspected or symptoms do not respond to conservative treatment.

### Pre-Op Care:

- Monitor for respiratory difficulty from inflammation or hematoma.
- Encourage coughing, deep breathing, and early ambulation.
- Monitor for hoarseness and inability to cough effectively.
- Assess the surgical dressing; monitor the surgical wound for infection, swelling, redness, drainage, or pain.
- **DIET:** Soft diet
- Monitor for sudden return of radicular pain; may indicate cervical spine instability.

## Post-Op Care:

- Assess the surgical dressing, bleeding, drainage, and surgical drains.
- Monitor lower extremities for sensation, movement, color, temperature, and paresthesia.
- Monitor for urinary retention, paralytic ileus, and constipation.
- **DIET:** High fiber diet; increase oral fluid intake (OFI)
- Administer opioids and sedatives as prescribed to relieve pain and anxiety.
- Assist regarding the use of back brace or corset and to wear cotton underwear to prevent skin irritation.
- **Position:** Lie in supine; place a pillow under the neck and slightly flex the knees.
- Avoid spinal flexion or twisting and that the spine should be kept aligned.
- Avoid extreme hip flexion when lying on the side.
- Following disk surgery, instruct the client in correct **logrolling techniques** for turning and repositioning and for getting out of bed.

**Amputation** is the surgical removal of a limb or part of the limb.

## Post-Op Care:

- Monitor for signs of complications (hemorrhage, infection, phantom limb pain, neuroma, and flexion contractures)
- Mark bleeding and drainage on the dressing if it occurs.
- Evaluate for phantom limb sensation and pain; medicate immediately.
- *Do not elevate the residual limb on a pillow.*
- **First 24 hours:** Elevate the foot of the bed to reduce edema; then keep the bed flat to prevent hip flexion contractures,
- **After 24 to 48 hours:** position the client prone to stretch the muscles and prevent hip flexion contractures.
- Maintain surgical application of dressing, elastic compression wrap, or elastic stump.
- *Massage the skin toward the suture line* to mobilize scar and prevent its adherence to underlying bone.
- Provide emotional support for the loss of body part.

## Interventions for below-knee amputation:

### " NAPE "

- **Not** to hang residual limbs over the edge of the bed.
- **Alignment:** Discourage long periods of sitting and knee flexion.
- **Prone** position
- **Edema** prevention

## Interventions for above-knee amputation:

### " 2Ps "

- **P**osition in prone
- **P**revent internal or external rotation of the limb; use sandbag, rolled towel, or trochanter roll along the outside of the thigh.

**Rheumatoid Arthritis (RA)** is a chronic systemic inflammatory disease (immune complex disorder); the cause may be related to a combination of environmental and genetic factors; leads to destruction of connective tissue and synovial membrane within the joints.

### Signs and Symptoms:

- Inflammation, tenderness, and stiffness of the joints
- Morning stiffness lasting longer than 30 minutes.
- Spongy, soft feeling in the joints
- Low-grade temperature, fatigue, and weakness
- Elevated ESR and positive rheumatoid factor (*Reference interval: Negative or < 60 units/mL*)
- Synovial tissue biopsy reveals inflammation

### Nursing Interventions:

- Provide range-of-motion exercise.
- Splints may be used during acute inflammation to prevent deformity.
- Apply heat or cold therapy as prescribed to joints.
- Avoid weight-bearing on inflamed joints.
- Identify and correct safety hazards in the home.
- Encourage the client to verbalize feelings.
- Assist the client with self-care activities and grooming.

**Osteoarthritis (OA)** is marked by progressive deterioration of the articular cartilage; causes bone buildup and the loss of articular cartilage in peripheral and axial joints.

### Signs and Symptoms:

- Joint pain that *diminishes after rest and intensifies after activity*.
- Pain occurs with slight motion or even at rest.
- Aggravated by temperature change and climate humidity.
- Difficulty getting up after prolonged sitting.
- Presence of **Heberden's nodes** (*high*) or **Bouchard's nodes** (*below*)

### Nursing Interventions:

- Administer medications as prescribed, such as acetaminophen or topical applications, NSAIDs and muscle relaxants.
- Position joints in function position and avoid flexion of knees and hips.
- Prepare the client for corticosteroid injections into joints as prescribed.

- Provide a bed or foot cradle to keep linen off feet and legs until inflammation subsides.
- Avoid large pillows under the head or knees.
- Immobilize the affected joint with a splint or brace until inflammation subsides.
- Apply **cold applications** as prescribed when the joint is acutely inflamed.
- Instruct the client to balance activity with rest and to participate in an exercise program.

**Gout** is a systemic disease in which urate crystals deposit in joints and other body tissues, results from abnormal amounts of uric acid in the body.

### Types:

- **Primary gout** results from a disorder of purine metabolism.
- **Secondary gout** involves excessive uric acid in the blood caused by another disease.

### Phases:

1. **Asymptomatic:** Client has no symptoms, but serum uric acid level is elevated.
2. **Acute:** Client has excruciating pain and inflammation of 1 or more small joints, especially the great toe.
3. **Intermittent:** Client has intermittent periods without symptoms between acute attacks.
4. **Chronic:** Results in deposits of urate crystals under the skin, within major organs, such as the kidneys, leading to organ dysfunction.

### Signs and Symptoms:

- Swelling and inflammation of the joints, leading to excruciating pain.
- **Tophi:** Hard, irregularly shaped nodules in the skin containing chalky deposits of sodium urate
- Low-grade fever, malaise, and headache.
- Pruritus from urate crystals in the skin
- Presence of renal stones from elevated uric acid levels

### Nursing Interventions:

- **DIET:** Low-purine diet; avoiding foods such as organ meats, wines, and aged cheese.
- Encourage a high fluid intake of 2000 mL/day.
- Encourage a weight reduction diet.
- Avoid alcohol and starvation diets because they may precipitate a gout attack.
- Increase urinary pH (above 6) by eating alkaline ash foods (i.e., green beans, broccoli).
- Monitor joint range-of-motion ability and appearance of joints.
- Provide heat or cold for local treatments to affected joint as prescribed.
- Administer medications such as analgesic, anti-inflammatory, and uricosuric agents as prescribed.
- Position the joint in mild flexion during acute attack.

**Osteoporosis** occurs post-menopausal or as a result of a metabolic disorder or calcium deficiency, most commonly in the wrist, hip, and vertebral column.

### Types:

## Primary osteoporosis

- Most often occurs in postmenopausal women; occurs in men with low testosterone levels
- Risk factors include decreased calcium intake, deficient estrogen, and sedentary lifestyle.

## Secondary osteoporosis

- Causes include prolonged therapy with corticosteroids, thyroid-reducing medications, aluminum-containing antacids, or antiseizure medications.
- Associated with immobility, alcoholism, malnutrition, or malabsorption.

## Risk Factors for Osteoporosis

? Cigarette smoking

? Early menopause

? Excessive use of alcohol

? Family history

? Female gender

? Increasing age

? Insufficient intake of calcium

? Sedentary lifestyle

? Thin, small frame

? White (European descent) or Asian race

## Signs and Symptoms:

- Back pain that occurs after lifting, bending, or stooping.
- Pelvic or hip pain, especially with weight-bearing.
- Decline in height from vertebral compression.
- *Kyphosis* of the dorsal spine, also known as “**dowager’s hump**”

## Nursing Interventions:

- Safety measures
- Clear walkway at home.
- Use side rails to prevent falls.
- Instruct in use of assistive devices such as a cane or walker.
- Use of a firm mattress.
- Move the client gently when turning and repositioning.
- Provide gentle range-of-motion exercises.

- Apply a back brace as prescribed during an acute phase.
- **DIET:** high in protein, calcium, vitamins C and D, and iron
- Maintain an *adequate fluid intake* to prevent renal calculi.